

**EFFECT OF DIFFERENT LEVEL OF NITROGEN FERTILIZER ON  
GROWTH AND YIELD AND YIELD COMPONENT OF TOMATO  
(*LYCOPERSICON ESCULENTUM* MILL.) AT WEST SHOWA ZONE,  
OROMIA, ETHIOPIA**

Nemomsa Beyene and Tesfaye Amare

Ambo University, College of Agriculture and Veterinary Sciences,  
Department of Plant Sciences, P. O. Box 19 Ambo, Ethiopia.

**ABSTRACT**

The field experiment was conducted in West showa zone, Toke kutaye district of Ormia region, Ethiopia with the objective to determine the optimum nitrogen fertilizer rate on different growth parameters, yield and yield component of tomato crop. To attain the objective, four level of nitrogen fertilizer Viz., 0, 50, 100, and 150kg/ha were used as treatments. The experiment was laid out in randomized complete block design (RCBD) with three replications. Data on plant height, primary branches per plant, number of leaves per plants, number of cluster per plant, number of fruits per cluster and fruit yield were collected from five plants of the middle rows of each plot and subjected to statistical analysis software (SAS) version 9.3 and LSD at 5% was used for mean comparison. The statistical analysis showed that, there was the significant ( $P < 0.05$ ) difference among treatments for all parameters except for number of fruit per cluster. For growth variables, 150kg/ha revealed the highest value but there was no significant ( $P < 0.05$ ) difference between 100 and 150kg/ha of nitrogen except for the height of the plant. The treatment 150kg/ha nitrogen fertilizer provided 22.41, 35.57 and 25.40% over the control treatment in height, number of leaves and primary branch, respectively. The treatment 150kg/ha nitrogen fertilizer increased the number of cluster per plant and yield of tomato fruit per hectare by 34.50 and 70.79% over the control treatment, respectively,. However, there was no significant difference in both number of cluster per plant and yield per hectare between 150 and 100ka N/ha.

**Keywords:** Tomato, nitrogen fertilizer, yield components, yield

**1. INTRODUCTION**

Tomato (*Lycopersicon esculentum mill*) belongs to the family Solanacea, genus *Lycopersicon*, sub family *Solanoideae* and tribe *Solaneae* [1]. It is widely cultivated in tropical, subtropical and

temperate climates and ranks third next to potato and sweet potato in terms of world vegetable production [2] with estimated annual area coverage and total production of 164 metric million tones which was harvested from 4.73 million hectares [3].

Among vegetable crops, tomato is the most important edible and nutritious worldwide [2]. Tomato plays an important role in human nutrition by providing essential amino acids, vitamins, minerals, sugars and dietary fibers [4]. The authors also reported that the fruit contains vitamin B and C, iron and phosphorus and its vitamin C content is particularly high and it is an important source of antioxidant such as lycopene. Tomato is almost always used as an essential ingredient in the diet of the people and in almost every household and usually used for preparing foods like soups, sauces, stews, salads and others, in cooked or processed form in large quantities as compared to other vegetables [5].

In Ethiopia, the highest share of commercial vegetable is taken by tomato [6]. Edossa *et al.* [7] indicated that the climatic and soil condition of Ethiopia allow cultivation of wide range of fruit and vegetable crops including tomato. The crop is produced at altitude range of 700 up to 2200 meter above sea level and with about 700 to over 1400 mm annual rain fall on different soils and different weather conditions in Ethiopia [8]. The average yield of tomato in Ethiopia is extremely low, 8 tone/ha compared to the world average yields which is 34 tone/ha [9]. Tomato production in Ethiopia reduced from 6298.63 ha area coverage and 283648.27 quintal yield in 2016/17 to 5235.19 ha and 277,745.38 quintals yield in 2017/18 cropping season [10].

The major yield reducing factors of tomato in Ethiopia are low soil fertility, erratic rainfall distribution, lack of high yielding cultivars, low soil moisture content, disease and insect pest and lack of optimum recommended fertilizer application rate [11]. Sanchez *et al.* [12] reported that, low fertilizer utilization and soil nutrient depletion is the major reason for low agricultural productivity in African country particularly in Ethiopia. Among these factors, application of optimum N fertilizer rate is one of the main determinant factors which significantly affect growth and yield of tomato in Ethiopia [13].

Tomato crop is highly responsive to nitrogen fertilizer application where nitrogen availability may be limited and time of the application is critical [14]. Hokam *et al.* [15] reported that nitrogen promotes vegetative growth and fruit yield, favors fruit development (when applied at later growing stage) and application of proper amount of the fertilizer has a dramatic effect on tomato growth and development. So far, fertilizer rates for tomato crop was determined only at Melkasa Research Center which cannot represent the agro-ecologically for other tomato growing regions of the country and especially no such study was done for tomato under vertisol condition [16]. Therefore, it is prudent to test different rate of nitrogen fertilizer at different topographic

area. Hence, the objective of this study was to determine the optimum nitrogen fertilizer level for growth, yield and yield components of Roma VF tomato variety at gudar area and for the areas of the similar soil type.

## **2. MATERIALS AND METHODS**

### **2.1. Description of the Study Area**

The study was conducted in West Showa zone, Toke kutaye district at Guder kebele, in 2019 under irrigation. The site is located at distance of 137 km to west of Addis Ababa (capital city) at 8°58'North latitude and 37°46'East longitude with an elevation 2101 meters above sea level. The annual range of temperature of the area is 15-40°C, with rain fall which ranges from 1257-2000mm. The soil types of the area is *vertisol* with pH value of 5.5-6.5.

### **2.2. Treatments and Experimental Design**

Roma VF tomato variety was used for the study. The seedling was prepared a head on a well prepared seed bed. The experiment was consisted of four nitrogen fertilizer rate (0, 50, 100, 150 kg/ha). The experiment was arranged in randomized complete block design (RCBD) with three replications. Each plot had 3.2 meter width and 2 meter length. The spacing between the blocks and plots were 1.5m and 1m, respectively. Four rows per plot with 0.8m and 0.4m between the row and the plant, respectively, was used. Each row contained five plants. The treatments were assigned to the plots randomly.

### **2.3. Experimental Procedure, Data collection and Data Analysis**

The seedlings were transplanted to the experimental field accordingly. At transplanting, 50% N from the total of each treatment was added to the experimental plot and remaining 50% of N was added after 45 days of transplanting. Five plants from the middle two rows of each plot were randomly taken and tagged for data collection. Plant height, number of leaf per plant, number of primary branches, number of cluster per plant, number of fruit per cluster and fruit yield per hectare data were collected.

The collected data were subjected to SAS software version 9.3 for Analysis of Variance (ANOVA) and means comparisons for the significantly different variables were made among treatments using Least Significant Differences (LSD) test at 0.05 levels of significance.

## **3. RESULT AND DISCUSSION**

### **3.1. Effect of nitrogen fertilizer on growth of tomato**

The statistical analysis showed that there was significant ( $P < 0.05$ ) difference among treatments in affecting the height of the plant. The longest plant (61.26cm) recorded from the plot treated by 150 kg N/ha and followed by (55.43cm) from the plot treated by 100 Kg N/ha. The shortest plant (47.53cm) recorded from the control plot (Table 1). The height of tomato plant was increased with the rate nitrogen fertilizer. However, there was no significant difference between 0 and 50kg N/ha in influencing the height of tomato. This result was in agreement with the result obtained by Biswas *et al.* [17] who revealed that the tallest plant was obtained by applying 150kg N/ha. This study also in line with the study conducted by Najafvand *et al.* [18] who reported that as the amount of nitrogen fertilizer increase the height of tomato also increase. Applying 150 KgN/ha increases plant height by 22.41, 19.03 and 9.5% compared to the control (0kgN/ha), 50, and 100kg/ha, respectively while, 100kg/ha increased plant height by 14.25% compared to the control which was in line with the study conducted by Gezu *et al.* [19] who revealed 13.6% in plant height increment compared to the control (no fertilizer application) by application of 99kg N/ha that application. The increase in plant height could be due to the readily available nitrogen which promotes vegetative growth and development. Nitrogen nutrient is responsible for photosynthesis, formation of chlorophyll and nucleic acids, its absence or deficiency causes stunted growth [20], hence this nutrient responsible for accumulation of greater biomass compared to unfertilized plot. These results was in in line with the results obtained by Akbar [21] who found that plant height in tomato increased with increase in nitrogen rate.

The study revealed that there was significant ( $P < 0.05$ ) difference between 0kg N/ha and 100kg N/ha, 0kg/h and 150kg/ha, 50kg/ha and 100kg/ha, and 50kg/ha and 150kg/ha in influencing the number of leaves per plant. The study revealed that, as the rate of nitrogen increase, the number of leaves per plant also increases. The highest leaf number per plant was recorded from the plot treated with 150kg N/ha. Biswas *et al.* [17] also reported the heights number of leaves per plant of tomato from the plot treated by 150kg N/ha. However, application of 100 kg N/ha was at par with 150kgN/ha on influencing number of leaves per plants and also using 50 kgN/ha also not different from using 0 kg N/ha on number of leaves per plant. Many authors reported that supplementary application of nitrogen fertilizer increase number of leaves per plant [21]. This result was in agreement with the result of Balemi [13] who reported that as nitrogen fertilizer level increases the number of tomato leaf increased.

Number of primary branches per plant was significantly ( $P < 0.05$ ) affected by rate of nitrogen fertilizer. The number of primary branch at 150 and 100 kg N/ha was 5.0 and 4.93, respectively, which was reduced to 4.08 and 3.73 when 50 and 0 kg N/ha applied, respectively. Application of 150kg/ha did not make significant difference in number of primary branch compared to 100kg/ha but significantly different from 50 and 0kg/ha. Iqbal *et al.* [22] also reported that application of 90 kg N/ha resulted in 4.33 primary branches which is in par using 100kg/ha in this study.

Application of 100kg N/ha increase the number of branch by 24.34% compared to application of 0kg nitrogen. The number of branch per plant increased with increasing nitrogen application up to optimum level. [19]also reported that, as the rate of nitrogen increases from 0-99kg/ha, the number branches in tomato increase and 99kg/ha nitrogen application increase number of branches by about 28.9% compared to the nil nitrogen.

**Table 1: Effect of nitrogen fertilizer rate on plant height, leaf number, branch number of tomato.**

Treatment N kg/ha	Plant height (cm)	Number of leaves per plant	Number of primary branch
0	47.53 <sup>c</sup>	27.53 <sup>b</sup>	3.73 <sup>c</sup>
50	49.60 <sup>c</sup>	31.20 <sup>b</sup>	4.08 <sup>c</sup>
100	55.43 <sup>b</sup>	38.60 <sup>a</sup>	4.93 <sup>ab</sup>
150	61.26 <sup>a</sup>	42.73 <sup>a</sup>	5.0 <sup>a</sup>
Mean	53.46	35.01	4.44
LSD (5%)	5.49	5.38	0.83
CV (%)	5.12	7.69	10.417

*Means within the same column followed by the same letter(s) are not significantly different from each other and different letter(s) indicated significantly different at 5% level of significance.*

### 3.2. Effect of nitrogen on yield component of tomato

Application of nitrogen fertilizer brought significant ( $P < 0.05$ ) change on number of cluster per plant of tomato. The highest number of cluster per plant was recorded from the plot received 150kg N/ha (20.40) followed by 100kg N/ha (19.26) which was significantly different from the plot treated by 50 and 0kg/ha (Table 2). However, there was no significant different between the plots treated by 100 and 150kg/ha in number of cluster per plant. In this study, as the nitrogen fertilizer level increased, the number of clusters per plant also increased however, the successive rates of fertilizer were not significantly different in affecting number of cluster per plant of tomato. The most probable reason could be tomato cluster formation do not give significant response for narrow rate of fertilizer. This result was in line with the result obtained by Iqbal *et al.*[22]. Biswas *et al.* [17] also reported highest number of cluster per plant of tomato from the plot treated by 150kg N/ha.

The analysis of the result demonstrated that the various doses of nitrogen fertilizer showed non-significant on number of fruits per cluster of tomato even if the mean values were different from

each other. This result was disagreed with the result revealed by Biswas *et al.* [17] who reported significant difference among nitrogen fertilizer levels in number of fruits per cluster but on different cultivar (BARI Tomato-9).

### 3.3. Effect of nitrogen on yield of tomato

The result of analysis showed that, there was significance ( $P < 0.05$ ) difference among treatments in influencing yield of tomato. The highest yield was obtained from the plot treated by 150 kg N/ha (28.737t/ha) which was not significantly different from the plot treated by 100kg N/ha (28.233t/ha). The lowest yield (8.393t/ha) was recorded from the plot treated by nil nitrogen fertilizer. There was significant differences between 0kg and 100kgN/ha and between 50 and 100kgN/ha in influencing the yield of tomato. This result revealed that the gradual increase in yield of tomato with the rate of nitrogen fertilizer up to optimum level. This result was in line with the finding of Biswas *et al.* [17] who reported that the highest fruit yield from the plot treated with 150kg N /ha. Application of 150 and 100kg N/ha increased yield of tomato by 70.79 and 70.27%, respectively, compared to the control treatment. Application of 50kgN/ha had 9.68t/ha yield advantage over the nil (control) fertilizer application. The result of this study was in line with result of Warner *et al.* [23] who reported that, as the rate of nitrogen fertilizer increases, the yield of tomato increase. Biswas *et al.* [17] also reported the highest fruit yield of tomato when the crop treated by 108.6 kg N /ha at the eastern part of Ethiopian country.

**Table 2: Effect of different rate of nitrogen fertilizer on number of clusters per plant, number of fruits per clusters and yield in tone per hectare of tomato.**

Treatment N kg/ha	Number of cluster/plant	Number of fruit/cluster	Yield t/ha
0	13.36 <sup>c</sup>	4.5333 <sup>a</sup>	8.393 <sup>c</sup>
50	14.20 <sup>bc</sup>	4.53 <sup>a</sup>	18.073 <sup>b</sup>
100	19.26 <sup>ab</sup>	4.80 <sup>a</sup>	28.233 <sup>a</sup>
150	20.40 <sup>a</sup>	5.06 <sup>a</sup>	28.737 <sup>a</sup>
Mean	16.88	4.73	20.867
LSD (5%)	5.72	0.81	2.1941
CV (%)	16.81	8.59	5.27

*Means within the same column followed by the same letter(s) are not significantly different from each other and different letter(s) indicated significantly different at 5% level of significance.*

#### **4. CONCLUSION**

Soil nutrient is one of important factor to attain optimum yield in all crop production system. Nitrogen nutrient is responsible for photosynthesis, formation of chlorophyll and nucleic acids and its absence or deficiency causes stunted growth. Hence, nitrogen plays important role in vegetative and fruit development in crops. Tomato (*Lycopersicon esculentum mill*) is third most important vegetable crop in the world. Tomato production is challenged by many factors from which optimum amount of nitrogen nutrient is the primary problem particularly in Ethiopia since the country characterized by diverse agro-ecological system.

Currently the farmers in the study area using the rate of nitrogen fertilizer that was determined by Melkasa Research Centre, which is about 300km far from the study area and with different soil type. Therefore, it was prudent to test different rate of nitrogen fertilizer to determine the optimum rate for tomato production in the study area and areas with similar soil type. To attain the objective, four levels of nitrogen fertilizer rate (0, 50, 100, and 150kg/ha) nitrogen fertilizer in the form of urea was used. The rate 100 and 150kg/ha provided 28.233 and 28.737 t/ha yield, respectively, in which there was no significant difference between the two treatment. Hence, the rate 100kg/ha nitrogen fertilizer can be recommended for optimum yield of tomato (Roma VF variety) in the study area and areas with similar soil type.

#### **REFERENCES**

- [1]. Jones G (1999). An approach to the genetics of nitrogen use efficient in Tomato. *Journal of experimental botany* 55:295-306.
- [2]. FAOSTAT (2010). Statistical data base of food and agriculture of the write donation Maria, Jose Diez and Fernano neuzh and book of plant breeding vegetable (152-156).
- [3]. FAO (2014) United Nation Food and Agriculture Organization Statistics Division. Crop Production data. Rome, Italy.
- [4]. Kanyomeka L and Shuvite B (2005). Influence of pruning on tomato production under controlled environments, *Agricultural Tropicaet Subtropica* 32(2):79-81
- [5]. Ellis J G (1998). Postharvest problems of tomato production in Ghana -Field studies of some selected major growing areas in Ghana. *Journal of the Ghana science association* 1(1):55-59.

- [6]. Tesfa B, Yosef A, Jibicho G, Gebeyehu W, Melkamu H (2016). Performance of introduced hybrid tomato (*Solanum lycopersicum* Mill.) cultivars in the Rift Valley, Ethiopia. *International Journal of Research in Agriculture and Forestry* 3(10): 25-28.
- [7]. Edossa E, Dechasa N, Alamirew T, Alemayehu Y, Desalegne L (2013b). Household fertilizer use and soil fertility management practices in vegetables crop production in the central rift valley of Ethiopia. *Science technol. Arts res. J.* 2(4) :47-55.
- [8]. Birhanu K, Tilahun K (2010). Fruit yield and quality of drip-irrigated tomato under deficit irrigation. *Afr. J. Food, Agric, Nutr. Dev.* 10 (2): 2139-2144.
- [9]. FAOSTAT data ( 2012). Agricultural data. Provisional 2012 Production Indices Data. Crop Primary. (<http://apps.fao.org/default.jsp>).
- [10]. CSA ( 2018). Report on area and production of major in Ethiopia. Central statistics Agency. Addis Abeba.
- [11]. Desalegne L (2002). Tomato research experience and production prospects, Ethiopian Agricultural Research Organization, 2002. *Research Report 43*. Addis Abeba, Ethiopia.
- [12]. Sanchez P A, Buresh R J, Izac A N, Place F M, Shepherd K D (1997). Soil fertility replenishment in Africa: SSSA Spec. Publ. 51. SSSA and ASA, Madison, WI. p. 1-46.
- [13]. Balemi T (2008). Response of tomato cultivars differing in growth habit to nitrogen and phosphorus fertilizers and spacing on vertisol in Ethiopia. *Acta Agric. Slov.* 91(1):103.
- [14]. Fontes P C R, Ronchi C P (2002). Critical values of nitrogen indices in tomato plants grown in soil and nutrient solution determined by different statistical procedures. *Pesq. agropec. bras., Brasília* 37(10): 1421-1429.
- [15]. Hokam E M, El -Hendawy S E , Schmidhalter U (2011). Drip Irrigation Frequency: The Effects and Their Interaction with Nitrogen Fertilization on Maize Growth and Nitrogen Use Efficiency under Arid Conditions. *J. Agronomy & Crop Science* 197: 186–201
- [16]. Kahsay Y K, Embaye A, Tekle G (2016). Determination of Optimum Rates of N and P Fertilizer for Tomato at Mereb-lekhe District, Northern Ethiopia. *Journal of Agriculture and Crops* 2(3): 24-30.



- [17]. Biswas M, Sarkar D R, Asif M I, Sikder R K, Mehraj H Jamal Uddin A F M (2015). Nitrogen Levels on Morphological and Yield Response of BARI Tomato - 9. *Journal of Science, Technology & Environment Informatics* 01(02): 68–74.
- [18]. Najafvand S, Direkvandi N, Alemzadeh A, Sedighie D F (2008). Effect of Different Levels of Nitrogen Fertilizer with Two Types of Bio-Fertilizers on Growth and Yield of Two Cultivars of Tomato (*Lycopersicon esculentum* Mill). *Asian Journal of Plant Sciences* 7(8): 757-761.
- [19]. Degefa G , Benti G, Jafar M, Tadesse F, Berhanu H (2019). Effects of Intra-Row Spacing and N Fertilizer Rates on Yield and Yield Components of Tomato (*Lycopersicon Esculentum* L.) at Harawe, Eastern Ethiopia. *Journal of Plant Sciences* 7(1):8-12.
- [20]. Tisdale S L, Nelson W L, Beaton J D, Havlin J L (2003). Soil Fertility and Fertilizers. 5th Edn., Prentice-Hall of India, New Delhi, India.
- [21]. Ewulo B S, Sanni K O, Adesina J M (2015). Response of tomato (*Lycopersicum esculentum* var. mill) to different levels of nitrogen and phosphorus fertilizer in south western nigeria. *International journal of applied and pure science and agriculture* 1(1): 2394-5532.
- [22]. Iqbal M, Niamatullah M, Yousaf I, Munir M, Khan M Z (2011). Effect of nitrogen and potassium on growth, economical yield and yield components of tomato. *Sarhad J. Agric.* 27(4): 545-548.
- [23]. Warner J, Zhang T Q, Hao X (2004). Effects of nitrogen fertilization on fruit yield and quality of processing tomatoes. *Canadian journal of plant science* 84: 865–871.